Gardening Notes, July 2025

Finally, the hazy, lazy days of summer have arrived – something many of us longed for during those chilly days last winter and even in May! The temperatures are warm, the days are long and hopefully the rain is ample! The heat and humidity have certainly returned too, so try to garden during the cooler temperatures of the early mornings and evenings. Keep records on daily temperatures and rainfall, since summers' heat and night-time temperatures impact a plant as much as winters' cold. Always remember to wear a big hat, apply ample sunscreen to exposed skin and drink plenty of water while you garden!

Things to do: Garden

- Weed and water your containers. The plants are now beginning to approach their mature size and it is important to keep them well fed and hydrated. Water as needed, which is usually once per day. However, if the pot is rootbound and temperatures hot, some containers should be watered twice daily.
- If you used slow-release fertilizer pellets in your containers, you may wish to consider a reapplication near months end. Most fertilizer pellets only last 2 or 3 months and the rate of release increases as the temperatures rise above 85°F. For heavy feeders, like *Brugmansia* (picture above) you should supplement the slow-release feed with liquid fertilizer since these plants are always hungry and the slow-release granules

do not release the nutrients at a sufficient rate to support blooming and growth.

• Some annuals, such as Coleus, Plectranthus and Persian Shield (Strobilanthes) benefit from an occasional pinching of the tips, allowing lower buds to break dormancy and grow (as seen at right for Persian Shield). This allows the plants to become denser and prevents them from becoming leggy or excessively large and overwhelming any neighboring plants.



- Most hanging baskets should receive a serious cutback in order to renew the plants for a late July through September display.
- Most hanging baskets also benefit from receiving a 50% dilute liquid fertilizer every other week with fertilizer.

- Irrigate perennial and annual gardens for long periods in the morning, ensuring a deep penetration of water into the soil should rainfall become less than ample.
- It is now ok to remove the old foliage of Narcissus as it turns brown during late June and July since next year's bulbs have now fully developed.
- Consider adding some graceful summer blooming nonhardy bulbs such as Abyssinian Gladiolus (*Gladiolus murielae*), pictured at right. Plants will bloom 8-10 weeks after planting.
- Reapply mulch to retain moisture should it have become thin or disturbed through replanting of annuals or perennials or from weeding. Mulch only needs to be



- a total of 2-3" thick including mulch from previous applications and new additions. Resist the temptation to use weed barrier under the mulch. It initially helps keep weeds down, but as the pores fill, water and air pass through more slowly. In addition, weeds often start to grow in compost on top of the barrier.
- For repeat blooming Shrub and Tea Roses, deadhead or remove the old flower blossoms, give them a light feeding and remove any leaves from the plant or on the ground that have blackspot, as this will help to reduce future outbreaks.
- Any Japanese Beetles on roses or other plants can be destroyed by flicking them into soapy water.
- Some perennials can be lightly trimmed such as the early blooming Salvias (Salvia nemorosa) while other plants can be cut back more heavily such as Catnip (Nepeta faassenii) and Ribbon Canary Grass (Phalaris arundinacea var. picta) to initiate new flowering stems or attractive foliage for late summer and fall. Nepeta 'Walkers Low' is photographed at right 2 weeks after being cut back, with an unpruned specimen behind. The trimmed plant is much denser and compact. Make certain you provide adequate irrigation if you cut the plants back hard and rain is lacking.
- Not all plants should be cut back hard. I prefer to leave the dried flower heads of Yarrow (*Achillea*) and Astilbe on the plant, as they look attractive come the fall and winter. The drawback is you may get unwanted seedlings.
- Early July is the latest time to pinch back unruly Asters and Chrysanthemums. Pinching any later will delay bloom time too far into the fall for the blooms to be effective. It is already too late to pinch overly exuberant Sedums like 'Autumn Joy' as they will be producing flowers by month's end.
- Weed! Even with the limited rainfall and warmer weather of June, there is an

endless supply of weeds! It is best to get them young before they become too noticeable and the root system established. Worse yet, try not to let them go to seed, as the average lifespan of a seed is 7 years which equates to 7 more years of weeding! Mulching heavy does not equate to less weeds either, and will lead to more issues in the future since excessive mulch is not healthy for root systems.

- Provide a light pruning and training of rampant growing Wisteria stems, stray stems of Clematis, and other vines. For Wisteria, I generally prune to the second or third compound leaflet along the viny new growth. A heavy pruning will only invigorate excessive vegetative growth and potentially less flower bud development. Do not prune off the spurs that produced flowers this past spring (the seed pods will still be attached), since they will yield flowers for next spring! Also, remove Wisteria vines running along the ground before they invade other areas!
- Train the growing stems of rampant annual vines like Firecracker Vine (pictured at right) or Cardinal Vine. Often the stems will start to twin around unintended neighboring shrubs rather than the intended support.
- Prune water sprouts and suckers on small trees and large shrubs before they become too large and potentially begin to deform the plant.
- With young ornamental and shade trees, you may wish to prune off or thin the lowest drooping branches if you have not already done so, as they will continue to hang lower until growth ceases towards midmonth. These low hanging

branches provide excessive shade for plantings beneath, block important views of the garden and make walking about difficult. Cut these branches back to – but not

into – the branch collar at the base of the branch.

Continue to visit local garden centers as new materials are continuing to arrive and you may see some summer bloomers like *Cephalanthus occidentalis* (Buttonbush, pictured above right) that you may have overlooked in May. Mid-summer sales maybe beginning as well!







• For the more adventurous, visit National Parks and study where our native flora grows! It is interesting to see what conditions the plants are growing under and what other plants are growing in the local plant community! I recently visited

Shenandoah National Park and was intrigued to find *Hydrangea* arborescens (Smooth Hydrangea pictured above) growing happily with *Rubus odoratus* (Flowering Raspberry pictured at right) on slopped terrain! The *Rubus* flower is seen in the closing image below and the plants are also thornless! The Hydrangeas have entirely fertile florets and lack the typical ring of showy sterile florets typical



of Lacecaps for sale at Garden Centers. More fertile florets for pollinators and still very attractive!

<u>Vegetables</u>

- Peas, radishes, lettuce, spinach and other cool season crops need to be removed, composted and replaced by beans, beets or any preferred 80-day to maturity crop. Thin carrots to roughly 1" apart and beets to 2" to allow room for proper root development. For beets, the greens can be used to garnish a salad!
- Come July, summer squash and cucumbers are beginning to come of size; harvest them small (6-8") and frequently to ensure continued yields and tenderness. If you planted potatoes, they can also be harvested starting in mid-July. A second crop of cucumbers can be planted in early July to extend production as those planted earlier that will begin to lose vigor come late August and September.
- When watering tomatoes, make certain that the fruit remains dry in order to
 - reduce the occurrence of blossom end rot. Keep the tomatoes staked (as seen at right). By positioning poles along a row of tomatoes and holding the tomatoes upright between two strings (marked by the arrows), the fruits can be kept off the ground and air can circulate through the plant. It reduces disease risks and the ripened fruit can be more easily seen.



- Of course, you may just have one or two tomato plants and a simpler staking method is desired. At Willowwood Arboretum several bamboo poles were fashioned around the plants to keep the plants growing upright while looking attractive at the same time (as pictured below)!
- Check parsley for Black Swallowtail Butterfly eggs or caterpillars. You may also wish to allow them to develop!

- If your Zucchini Squash was infected with borers early this season and have collapsed, consider planting some of the Scallop Squash, often called Patty Pan Squash for an autumn harvest. They mature in about 50-54 days from seed and come in solid White, Golden, and a combination of White and Green Striped. Not only are they delicious, they will add great color to the Thanksgiving Table!
- Maintain even moisture in the soil around tomato plants to reduce the risk of the tomatoes splitting. This is particularly true of Heirloom tomatoes and using a straw mulch will help maintain consistent moisture.
- Yellowing of the lowest leaves on the stems of tomatoes is often an indication of Early Blight. It is best to remove these lower leaves, such that there is no contact with the soil and reducing the threat of future outbreaks. Also, wash your hands after removing the leaves before touching other plants to reduce the chance of spreading the disease.

Houseplants

- Move some of your favorites outdoors where possible. They will love the humidity, but remember to move them into a shaded location at the start! Many succulents like *Haworthiopsis* will develop red hues when moved outside and into the sun!
- If spring was too hectic and you are just getting caught up, July is a good time to up-pot those houseplants that have outgrown their container. Remember, pick a container that is roughly 2" bigger than the initial container, allowing an area of about 1" around the root ball for new roots to grow. Also, position the plant ½-1" below the rim of the new pot, allowing space for the irrigating water and preventing it from spilling over the sides.
- If you have Air Plants (*Tillandsia* spp.) or other epiphytic plants such as the *Neoregelia* 'Tiger Cub', tie them into trees with twine and replace the plastic pot with a burlap wrap. It is amazing how much they grow and produce pups with a minimal care!
- Houseplants that remain indoors are actively growing during the summer and should receive a regular watering regiment of every two weeks or so and monthly fertilizing with a 50% dilute liquid solution.

<u>Turf</u>

- Raise the cutting height to 3" during the hot and dry months, reducing plant stress.
- If you irrigate, do so for extended periods during the early morning to encourage deeper root growth. Try not to water in the evening, since that will promote various fungal problems.
- Make certain you check and adjust the orientation of sprinklers. I am amazed at the number of sprinklers I see irrigating roadways and not the intended turf!
- Avoid fertilizing during July through mid-August as that will necessitate more irrigation.


• Consider reducing the size of turf areas if the size is no longer needed. Using groundcover masses benefit native insects and reduce the negative impacts from lawnmowers, unnecessary irrigation and fertilizers.

Bulbs

- Although this maybe the time of year when bulbs are farthest from your mind,
 - bulb catalogues have been arriving now for the past couple of months and it is not too early to order! In fact, some bulbs are already sold out, so get those orders in.
- Lycoris squamigera, commonly called Surprise Lily or Naked Lady due to their ability to bloom unexpectedly and without foliage (pictured above right), will begin to bloom at the end of July. They



are tolerant of full sun or light shade and are a bulb that many gardeners do not think about since they bloom outside the normal bulb bloom time. Easily grown, consider adding this beauty to your garden later this summer or late next spring!





Rubus odoratus

Bruce Crawford Manager of Horticulture, Morris County Parks Commission